

## Imid-Triadimenol

Similar to: *Zorro*<sup>®</sup>

<b>Active Constituent</b> Imidacloprid 180g/L and Triadimenol 56g/L		<b>Formulation</b> SC	
<b>Typical Seed</b> Wheat, oats and barley (see registration section).			
<b>Pests Controlled</b> Wheat and Corn Aphid, Barley Yellow Dwarf Virus, Bunt, Powdery Mildew and Smuts, and suppression of Septoria Tritici Blotch, Stripe Rust and Leaf Scald.			
<b>Chemical Group</b> 4A Insecticide 3 Fungicide		<b>Mode of Action</b> Root uptake, upwardly systemic in plants, contact and ingestion activity on insects, and protectant, curative and eradicated activity on fungi.	
<b>Price ex GST</b> \$20.00/L		<b>Typical Pack Size</b> 10L	
<b>Poison Schedule</b> 5	<b>Dangerous Goods Class</b> -	<b>UN</b> -	<b>Packing Code</b> -
<b>Withholding Period</b>			
<b>Harvest</b> Not required when used as directed.			
<b>Grazing</b> Do not graze or cut for stock food for 9 weeks post sowing.			
<b>Application/Efficacy</b> Agitate container vigorously to ensure product is thoroughly suspended prior to applying or adding to water. Apply a maximum of 600ml of mixture to 100kg of seed, ensuring thorough coverage. Calibrate seeding equipment with treated grain – flow rates may differ from untreated seed. Sowing depth should not exceed 50mm, and sowing rates may need to be increased to compensate for delayed crop emergence.			
<b>Treated Grain</b> Grain must be used within 12 months of treating.			
<b>Storage Life</b>			
<b>Compatibility</b> Do not mix with other products.			
<b>Incompatibility</b> N/A			
<b>4F Broadacre Registrations</b> Wheat, oats and barley (extract listed).			
<b>4F Other Registrations</b> N/A			
<b>Similar Product Registrations</b> <i>Zorro</i> <sup>®</sup> is also registered as a stored cereal grain protectant.			
<b>Situation</b>	<b>Target Pest/Disease</b>	<b>Rate/100kg seed</b>	<b>Comments</b>
Cereals	Wheat Aphid, Corn Aphid, BYDV	400ml	Synthetic Pyrethroid top-up spray should be applied 7-8 weeks post-sowing.
Wheat	Bunt, Smuts, SPP Stripe Rust, Septoria Tritici Blotch		
Oats	Smuts		
Barley	Smuts, Powdery Mildew, suppression of Leaf Scald		
<b>Comments</b>			
<b>Do not</b> allow grain intended for human or animal consumption to become contaminated with this product.			
<b>Do not</b> use treated seed for human or animal consumption.			
<b>Do not</b> apply to seed with moisture content higher than 12%.			
Imidacloprid can provide a certain level of crop abiotic stress protection in some situations.			