

Procymidone 500 SD

Similar to: *Sumisclex® Broadacre*

Active Constituent	Procymidone 500g/L		Formulation	SC
Typical Seed	Lupins (see registration section).			
Pests Controlled	Brown Leaf Spot.			
Chemical Group	2 Fungicide	Mode of Action	Upwardly systemic activity in plant – provides protectant, curative and eradicator activity on fungi.	
Price ex GST	\$15.00/L	Typical Pack Size	20L	
Poison Schedule	7	Dangerous Goods Class	-	UN - Packing Code -
Withholding Period	Harvest	Not required when used as directed.		
	Grazing	Do not graze or cut for stock food for 13 weeks post sowing.		
Application/Efficacy	Agitate container vigorously to ensure product is thoroughly suspended prior to adding to water. Apply 400ml of mixture to 100kg of seed, ensuring thorough coverage. High rates will reduce Rhizobium inoculum efficacy, however will not impact established soil Rhizobia populations. Do not use alkaline water, and persistence will be reduced in alkaline soils.			
Treated Grain	Grain must be used within 12 months of treating.			
Storage Life				
Compatibility	Not generally required.			
Incompatibility	N/A			
4F Broadacre Registrations	Lupins (extract listed).			
4F Other Registrations	N/A			
Similar Product Registrations	As above.			
Situation	Target Pest/Disease	Rate/100kg seed	Comments	
Lupins	Brown Leaf Spot	50-200mL	Use higher rate when disease risk and conditions are high.	
Comments				
Do not allow grain intended for human or animal consumption to become contaminated with this product.				
Do not use treated seed for human or animal consumption.				
Do not sow seed in overly wet and cold conditions, or areas prone to water logging.				
Known to cause birth defects in laboratory animals; women of childbearing age should avoid any contact.				
Birds and animals eating treated seed may be poisoned.				
Dangerous to aquatic animals.				
S7 product – TOXIC				