

## Tebuconazole 25T

Similar to: *Raxil*<sup>®</sup>

<b>Active Constituent</b>	Tebuconazole 25g/L and Triflumuron 4g/L		<b>Formulation</b> SC
<b>Typical Seed</b>	Wheat, barley and oats (see registration section).		
<b>Pests Controlled</b>	Bunt, Smuts (Flag and Loose), and Granary Weevil, Lesser Grain Borer, Rice Weevil, Rust-red Flour Beetle and Sawtoothed Grain Beetle.		
<b>Chemical Group</b>	3 Fungicide 15A Insecticide	<b>Mode of Action</b>	Upwardly systemic in plant – provides protectant, curative and eradicated activity on fungi, and contact and residual activity on insects.
<b>Price ex GST</b>	\$20.00/L	<b>Typical Pack Size</b>	10L
<b>Poison Schedule</b>	5	<b>Dangerous Goods Class</b>	-
		<b>UN</b>	-
		<b>Packing Code</b>	-
<b>Withholding Period</b>	<p><b>Harvest</b> Not required when used as directed.</p> <p><b>Grazing</b> Do not graze plants grown from treated seed or cut for stock food within 4 weeks of sowing.</p>		
<b>Application/Efficacy</b>	Agitate container vigorously to ensure product is thoroughly suspended prior to addition into clean water to form a cream. Add cream to required water volume in pot/tank and maintain agitation throughout mixing and application. Sowing depth should not exceed 50mm, and sowing rates may need to be increased to compensate for delayed crop emergence. For post-harvest treatment, apply to seed with moisture content below 12% to ensure adequate storage life.		
<b>Treated Grain Storage Life</b>	Use treated grain within 12 months.		
<b>Compatibility</b>	Not recommended to be mixed with other products (perform a jar test if required).		
<b>Incompatibility</b>	Do not mix with trace elements.		
<b>4F Broadacre Registrations</b>	Wheat, barley and oats		
<b>4F Other Registrations</b>	N/A		
<b>Similar Product Registrations</b>	As above.		
<b>Situation</b>	<b>Target Pest/Disease</b>	<b>Rate/100kg seed</b>	<b>Comments</b>
Wheat	Bunt, Flag Smut, Loose Smut	100mL	Mix with water for good coverage of seed. Apply 400ml total mixture per 100Kg of seed (do not exceed 600mL).
Barley, Oats	Covered and Loose Smut		
Cereals	Protection against pests of stored grains: Granary Weevil, Lesser Grain Borer, Rice Weevil, Rust-red Flour Beetle, Sawtoothed Grain Beetle		
<b>Comments</b>			
Do not allow grain intended for human or animal consumption to become contaminated with this product.			
Do not use treated seed for human or animal consumption.			
Treated seed is dangerous to birds and animals.			
Heavier soils prone to crusting or clodding may cause delayed crop emergence, particularly in cold conditions.			
Perform a germination test on treated seed that is stored in excess of the 12 months.			