

Triadimenol 150+ (Powder)

Similar to: *Baytan C®*

Active Constituent	Triadimenol 150g/kg and Cypermethrin 4g/kg		Formulation	WP
Typical Seed	Wheat, barley and oats.			
Pests Controlled	Bunt, Leaf Scald, Powdery Mildew, Smuts, Stripe Rust, and Granary Weevil, Lesser Grain Borer, Rice Weevil, Rust-red Flour Beetle and Sawtoothed Grain Beetle.			
Chemical Group	3 Fungicide 3A Insecticide	Mode of Action	Fungicide is upwardly systemic in plant – provides protectant, curative and eradivative activity on fungi. Insecticide provides ingestion, contact and residual activity on insects.	
Price ex GST	\$18.00/kg	Typical Pack Size	10kg	
Poison Schedule	5	Dangerous Goods Class	-	UN - Packing Code -
Withholding Period	Harvest Not required when used as directed. Grazing Do not graze or cut for stock food for 5 weeks post sowing.			
Application/Efficacy	Agitate container vigorously to ensure product is thoroughly suspended prior to addition into clean water to form a cream. Add cream to required water volume in pot/tank and maintain agitation throughout mixing and application. Sowing depth should not exceed 50mm, and sowing rates may need to be increased to compensate for delayed crop emergence. For post-harvest treatment, apply to seed with moisture content below 12% to ensure adequate storage life.			
Treated Grain Storage Life	Use treated grain within 12 months.			
Compatibility	Not recommended to be mixed with other products (perform a jar test if required).			
Incompatibility	Do not mix with trace elements, and do not use alkaline water.			
4F Broadacre Registrations	Wheat, barley and oats.			
4F Other Registrations	N/A			
Similar Product Registrations	As above.			
Situation	Target Pest/Disease	Rate/100kg seed	Comments	
Wheat	Bunt, Flag Smuts, suppression of Stripe Rust	100-150g	Use in a total mixture of 400-600ml/100kg seed.	
Oats and Barley	Loose and Covered Smut, suppression of Powdery Mildew and Leaf Scald		Use the higher rate when foliar diseases more frequently appear.	
Wheat, Oats and Barley	Granary Weevil, Indian Meal Moth, Lesser Grain Borer, Rice Weevil, Rust-red Flour Beetle, Sawtoothed Grain Beetle, Tropical Warehouse Moth			
Comments				
<p>Do not allow grain intended for human or animal consumption to become contaminated with this product.</p> <p>Do not use treated seed for human or animal consumption.</p> <p>Also available in a soluble concentrate solution.</p> <p>Contains blue dye for identification.</p> <p>Treated seed is toxic to birds and animals.</p> <p>Heavier soils prone to crusting or clodding may cause delayed crop emergence.</p> <p>Perform a germination test on treated seed that is stored in excess of the 12 months.</p>				