

Triticonazole 200 C

Similar to: *Real 200*[®]

Active Constituent	Triticonazole 200g/L and Cypermethrin 4g/L		Formulation	SC
Typical Seed	Wheat and barley.			
Pests Controlled	Smuts, and suppression of Leaf Rust, Stripe Rust, Powdery Mildew and Leaf Scald, and Lesser Grain Borer, Granary Weevil, Rice Weevil, Rust-red Flour Beetle, Saw Toothed Grain Beetle and Tropical Warehouse Moth.			
Chemical Group	3 Fungicide 3A Insecticide	Mode of Action	Fungicide is upwardly systemic in plant – provides protectant, curative and eradicated activity on fungi. Insecticide provides ingestion, contact and residual activity on insects.	
Price ex GST	\$80.00/L	Typical Pack Size	5L	
Poison Schedule	5	Dangerous Goods Class	-	UN - Packing Code -
Withholding Period	Harvest Not required when used as directed. Grazing Not required when used as directed.			
Application/Efficacy	Agitate container vigorously to ensure product is thoroughly suspended prior to addition into clean water. Add a minimum of 400ml of mixture (water + product) for every 100kg seed, ensuring thorough coverage.			
Treated Grain Storage Life	Use treated grain within 12 months.			
Compatibility	Do not mix with other products.			
Incompatibility	Do not mix with trace elements, and do not use alkaline water.			
4F Broadacre Registrations	Wheat and barley (extract listed).			
4F Other Registrations	N/A			
Similar Product Registrations	<i>Premis Pro C</i> [®] ($\frac{1}{8}$ strength of 4Farmers Triticonazole 200 C) is registered for cereal smuts at 100ml/100kg of seed.			
Situation	Target Pest/Disease	Rate/100kg seed	Comments	
Wheat	Covered Smut, Flag Smut, Loose Smut, suppression of Leaf Rust, Stripe Rust	75-150mL	Use higher rates where foliar diseases are more common, conditions favour the disease or when a prolonged protection is required.	
Barley	Covered and Loose Smut, suppression of Powdery Mildew, Leaf Scald			
Wheat, Barley	Lesser Grain Borer, Granary Weevil, Rice Weevil, Rust-red Flour Beetle, Saw Toothed Grain Beetle, Tropical Warehouse Moth			
Comments				
<p>Do not allow grain intended for human or animal consumption to become contaminated with this product.</p> <p>Do not use treated seed for human or animal consumption.</p> <p>Trials indicate Triticonazole does not reduce seedling coleoptile length.</p> <p>Triticonazole 200 C represents excellent value for the control of seed borne diseases and stored grain pests (see Registration Section comments).</p>				