

## Omethoate 290

Similar to: *Le-mat*<sup>®</sup>

<b>Active Constituent</b>	Omethoate 290g/L		<b>Formulation</b> SL	
<b>Typical Situations</b>	Barrier spray (firebreaks and fence lines).			
<b>Insects Controlled</b>	Red Legged Earth Mite.			
<b>Chemical Group</b>	1B	<b>Mode of Action</b>	Short-term systemic in plants, contact and ingestion with some surface residual activity.	
<b>Price ex GST</b>	\$30.00/L	<b>Typical Pack Size</b>	10L	
<b>Poison Schedule</b>	6	<b>Dangerous Goods Class</b>	6.1/3	<b>UN</b> 3017 <b>Packing Code</b> III
<b>Withholding Period</b>	<b>Harvest</b>	Not required when used as directed.		
	<b>Grazing</b>	Not required when used as directed.		
<b>Application Method</b>	Boom spray or offset jet 50-100L/ha		<b>Rain fastness</b>	1 hour
<b>Efficacy</b>	Ensure thorough coverage of both soil and weeds in the barrier area (via appropriate jet size and water rates).			
<b>Compatibility</b>	Compatible with most commonly used herbicides (conduct a jar test if unsure).			
<b>Incompatibility</b>	Not compatible with alkaline mixtures (e.g. Bordeaux mixture and lime).			
<b>Water Quality</b>	N/A			
<b>4F Broadacre Registrations</b>	N/A			
<b>4F Other Registrations</b>	Barrier spray (firebreaks and fence lines).			
<b>Similar Product Registrations</b>	As above.			
<b>Situation</b>	<b>Target Insect/s</b>	<b>Rate/ha</b>	<b>Comments</b>	
Barriers	Red Legged Earth Mite	300mL	To prevent an invasion, spray bare soil around the crop/pasture along the fence line.	
<b>Comments</b>				
<b>CARE:</b> Damaging to bees - to minimise impact on foraging bees, apply early morning or late evening. Omethoate can provide a residual of 1-2 weeks for the control of Red Legged Earth Mite.				