

## Epoxiconazole 125

Similar to: *Opus*<sup>®</sup>

<b>Active Constituent</b>	Epoxiconazole 125g/L		<b>Formulation</b>	SC
<b>Typical Situations</b>	Wheat and barley (see registration section).			
<b>Diseases Controlled</b>	Leaf Scald, Net Form Net Blotch, Powdery Mildew (Wheat & Barley), Rust (Leaf & Stripe) and Septoria Nodorum Blotch.			
<b>Chemical Group</b>	3	<b>Mode of Action</b>	Upwardly systemic in plant – provides protectant, curative and eradivative activity.	
<b>Price ex GST</b>	\$35.00/L	<b>Typical Pack Size</b>	20L	
<b>Poison Schedule</b>	5	<b>Dangerous Goods Class</b>	-	<b>UN</b> -
<b>Withholding Period</b>	<b>Harvest</b> Do not harvest for 6 weeks post application. <b>Grazing</b> Do not graze or cut for stock food for 6 weeks post application.			
<b>Application Method</b>	Boom spray 50-100L/ha, aerially at least 20L/ha		<b>Rain fastness</b>	When dry
<b>Efficacy</b>	Apply at recommended timing for specific situation, ideally in fine weather, ensuring thorough canopy penetration and foliage coverage to maximise efficacy. Fungicidal activity only occurs within treated foliage – new foliage is not protected.			
<b>Adjuvants</b>	Non-ionic wetters may assist coverage and efficacy where plants are difficult to wet.			
<b>Compatibility</b>	N/A			
<b>Incompatibility</b>	N/A			
<b>Water Quality</b>	N/A			
<b>4F Broadacre Registrations</b>	Wheat and barley (extract listed).			
<b>4F Other Registrations</b>	N/A			
<b>Similar Product Registrations</b>	As above.			
<b>Situation</b>	<b>Target Disease/s</b>	<b>Rate/ha</b>	<b>Comments</b>	
Wheat (Z32-59)	Leaf Rust	500mL	Apply when conditions favour disease development, prior to development in the crop. Re-spray may be required if infection occurs early; monitor crop for 3-4 weeks following applications. Use higher rate when disease is present on the top leaf.	
	Stripe Rust, Septoria Nodorum	250-500mL		
	Powdery Mildew	250mL		
Barley (Z32-59)	Leaf Rust	250-500mL	<b>Do not exceed 500mL/ha of Epoxiconazole 125 in any single growing season.</b> Apply when conditions favour disease development, prior to development in the crop. Re-spray may be required if infection occurs early; monitor crop for 3-4 weeks following applications. 2 applications at a lower rate may be more effective in some cases.	
	Net Form of Net Blotch	250-500mL		
	Leaf Scald, Powdery Mildew	250mL		
<b>Comments</b>				
<b>Do not</b> apply after the Z59 crop stage. Excellent product for Septoria and Rust control in cereals. Possesses excellent mixing properties.				