

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



TRIADIMENOL 250 E.C. Fungicide

ACTIVE CONSTITUENT: 250 g/L TRIADIMENOL
SOLVENT: 600 g/L N-METHYL-2-PYRROLIDONE

GROUP C FUNGICIDE

Cabbage; Cauliflower & Broccoli: Systemic control of Ring Spot.
Cucurbits; Grapevines; Papaws: For control of powdery mildew.
Sugar Cane: Control of Pineapple disease.

DO NOT DESTROY CONTAINER WHILE PRODUCT STILL REMAINS

READ THE ATTACHED INFORMATION THOROUGHLY BEFORE USING THIS PRODUCT

NET CONTENTS
 1L
 5L
 20L

APVMA Approval No.
 52067/1L/0102;
 52067/5L/0102;
 52067/20L/0102.

4FARMERS PTY. LTD.
 A.C.N. 067 443 485
 1/70 McDOWELL ST, WELSHPOOL,
 WESTERN AUSTRALIA 6106
 PH: (08) 9356 3445 FAX: (08) 9356 3447

GENERAL INSTRUCTIONS

4Farmers Triadimenol 250 EC Fungicide is a broad spectrum, systemic, foliar fungicide. As it is quickly taken up by plants and rapidly translocated with systemic activity, 4Farmers Triadimenol 250 EC Fungicide is less affected by rainfall or irrigation than contact type fungicides.

FUNGICIDE RESISTANCE WARNING

GROUP C FUNGICIDE

Triadimenol is a member of the D.M.I. group of fungicides. For fungicide resistance management, 4Farmers Triadimenol 250 is a Group C fungicide. Some naturally occurring individual fungi resistant to 4Farmers Triadimenol 250, and other Group C fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly. These resistant fungi will not be controlled by 4Farmers Triadimenol 250 and other Group C fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, 4Farmers Pty Ltd accepts no liability for any losses that may result from the failure of 4Farmers Triadimenol 250 to control resistant fungi.

MIXING

• Half fill the spray tank with clean water, then add the required quantity of 4Farmers Triadimenol 250 EC Fungicide with the agitator going. Fill to

top with water and agitate thoroughly.

- DO NOT tank mix with trace elements or insecticides without first checking compatibility.

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT

- Harmful to fish.
- DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

- Keep out of reach of children.
- Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to tank. Do not dispose of chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.
- If not recycling, puncture top, bottom and sides of empty container then crush and bury in an approved local authority landfill. If no landfill available, bury under at least 500 mm of soil in a disposal pit set up for this purpose, clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS:

Will irritate the eyes and skin. Avoid contact with eyes and skin. Do not inhale spray mist. When preparing the spray, wear cotton overalls buttoned to the neck and wrist, washable hat and elbow-length butyl rubber gloves and goggles. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID:

- If poisoning occurs, contact a doctor or Poisons Information Centre on 13 11 26 (24hrs).

MSDS:

For further information see Material Safety Data Sheet which can be obtained from the supplier.

CONDITIONS OF SALE:

The use of 4Farmers Triadimenol 250 EC Fungicide being beyond the control of the manufacturer, no warranty expressed or implied is given by 4Farmers Pty. Ltd. regarding its suitability, fitness or efficacy for any purpose for which it is used by the buyer, whether in accordance with the directions or not and 4Farmers Pty. Ltd. accepts no responsibility for any consequences whatsoever resulting from the use of this product.

| CROP | DISEASE/PEST | STATE | RATE | CRITICAL COMMENTS |
|------------------------------------|-------------------|----------------------|---|--|
| Cabbage Cauliflower Broccoli | Ring Spot | All States | 200 or 400 mL/ha Plus Agridex® 200 mL/ha | Apply every 10 days. Start the program immediately following the first two consecutive days after transplanting. Use the higher rate under heavy disease conditions favour development or the disease. |
| Cucurbits | Powdery Mildew | | 40 mL/100 L or 400 mL/ha | Apply a non-DMI fungicide such as Morestan® on a schedule commencing as soon as possible after emergence until fruit set. Then either tank mix or alternate Triadimenol with a non-DMI fungicide such as Morestan® on a 5-10 day schedule. Use Short Spray intervals when disease pressure is high or plants are growing rapidly. Good coverage is essential for good results. |
| Grapevines | | | 10 mL/100 L or 100 mL/ha | Apply thoroughly in the following spray program: 1. When shoots are 10 cm long. 2. Pre-flowering. 3. At flowering. 4. After fruit set. 5. Before bunches close. Do not apply in volumes less than 250 L/ha. For spray volumes between 250 and 1000 L/ha, use Triadimenol at 100 mL/ha. For volumes greater than 1000 L/ha, use the volume rate of 10 mL/100 L. |
| Papaws | | QLD, NSW, WA Only | 20 mL/100 L or 200 mL/ha | Spray when powdery mildew infection begins or when conditions favour the disease. Repeat application every 14 days while conditions favour the disease. Ensure thorough coverage of foliage particularly the crown region. |
| Sugar Cane | Pineapple Disease | QLD, NSW Only | 20 mL/100 L | Apply to setts by dipping or spraying. Ensure thorough wetting of cut ends. |

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:

CUCURBITS: Do Not harvest for 1 day after application.
CABBAGE, CAULIFLOWER, BROCCOLI, GRAPEVINES, PAPAWS: Do Not harvest for 7 days after application.
CABBAGE, CAULIFLOWER, BROCCOLI: Do Not graze any treated area, or cut for stock food.
SUGAR CANE: Not required when used as directed.

Batch Number:

Date of Manufacture:

TRIADIMENOL 250 E.C. Fungicide