

## CAUTION

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



# TRI-ALLATE 500 EC

Selective Herbicide

ACTIVE CONSTITUENT: 500 g/L TRI-ALLATE  
SOLVENT: 459 g/L LIQUID HYDROCARBON

**GROUP J HERBICIDE**

For the control of Wild oats in Wheat, Triticale, Chickpeas, Barley, Peas, Linseed, Lupins, Canola (Rapeseed), Faba beans and Safflower and for use in mixtures with 4Farmers Trifluralin 480 EC for control of a range of grass and broadleaf weeds in no-till/min-till cropping systems, pre-sowing or incorporated by sowing (IBS) as per directions for use.

**READ THIS LEAFLET BEFORE USING THIS PRODUCT**

**4 Farmers Australia Pty Ltd**

A.B.N. 51 160 092 428

1 / 70 McDowell Street, Welshpool, WA 6106

TEL: (08) 9356 3445 www.4farmers.com.au

Batch Number:

Date of Manufacture:

APVMA Approval No: 62333/56604

## DIRECTIONS FOR USE

Crop	Weed Controlled	States	Rate L/ha	Critical Comments
Barley, Triticale, Wheat, Chickpeas, Faba beans, Lupins, Peas, Linseed, Canola, Safflower	Wild Oats	All States	1.6 L	Apply immediately prior to or up to 3 weeks before sowing to a friable seedbed under conditions which will allow for the incorporation of the product to a depth of 2 to 4 cm with either a single or double pass of suitable Incorporation equipment (see Incorporation). A delay of 1 week between application and planting is recommended for Linseed. COMPLETE AND UNIFORM INCORPORATION OF 4Farmers TRI-ALLATE 500 EC Selective Herbicide IS ESSENTIAL FOR WEED CONTROL. See GENERAL INSTRUCTIONS for details of incorporation. 4Farmers TRI-ALLATE 500 EC Selective Herbicide may be used in crops undersown with legumes. For shallow-sown legume and oil seed crops, complete incorporation by cultivation to a depth of 5 cm must be undertaken before planting. 4Farmers TRI-ALLATE 500 EC Selective Herbicide may be tank mixed with 4Farmers CHLORSULFURON 750WDG for control of annual ryegrass, paradoxa grass and certain broadleaf weeds prior to planting of wheat only. 4Farmers TRI-ALLATE 500 EC Selective Herbicide may be tank mixed with 4Farmers TRIFLURALIN 480 EC for control of annual ryegrass, annual phalaris, paradoxa grass, enhanced control of wild oats (emerging from top 1 cm of seedbed) and wireweed. Do not apply less than 1 week prior to planting of wheat, barley, triticale and linseed (do not use the tank mix in Qld). Incorporation of this tank mix should be within 4-5 hours of application. Refer to the 4Farmers TRIFLURALIN 480 EC label incorporation details. Read and follow label instructions, restraints, plant-back periods and safety directions for tank mix products.
Wheat, barley and triticale	Annual Ryegrass, Wireweed, Phalaris spp, Fumitory, Wild Oats, Cereal oats, Sand Fescue ( <i>Vulpia fasciculata</i> ), Silvergrass ( <i>Vulpia bromoides</i> ) Wintergrass ( <i>Poa annua</i> ), Paradoxa Grass (Canary Grass) ( <i>Phalaris paradoxa</i> ), Corn Gromwell (Sheepweed) ( <i>Buglossoides anvensis</i> ), Rough Poppy ( <i>Papaver hybridum</i> )  Suppression Soil Surface – Bromegrass ( <i>Bromus diandrus</i> ), Barleygrass ( <i>Hordeum leporinum</i> ), Three Cornered Jack (doublegee) ( <i>Emex australis</i> ), Caltrop (Yellowvine & Bullhead) ( <i>Tribulus terrestris</i> ), Yellow Burr Weed ( <i>Amsinckia</i> spp), Deadnettle ( <i>Lamium amplexicaule</i> ), Speedwell ( <i>Veronica</i> spp)	WA, SA, Vic, NSW, Qld only	1.6-2.0L plus 1.5-2.0L 4Farmers TRIFLURALIN 480 EC	<b>For use in no-till/min-till Cropping Systems, Pre-Sowing or Incorporated by Sowing (IBS).</b> Use only with knife/blade points and presswheels – refer to table 13 for method of incorporation on the 4Farmers Trifluralin 480. Use higher rates on light sandy and sandy loam soils. <b>DO NOT use on heavy soils.</b> Avoid soils, which are non-wetting or are likely to become clumpy or cloddy as they may suffer reduced weed control. Use higher rates for heavier stubble coverage and high weed density situations. Stubble coverage above 40-50% ground cover can reduce weed control below acceptable levels. Suppression of Bromegrass and Barleygrass may be reduced in medium to high weed density situations. Higher rates are likely to provide improved control levels of Bromegrass, Wild oats, Cereal oats, Barleygrass and Silvergrass. Control of deep/late germinating weeds may be reduced. Insufficient incorporation of 4Farmers TRI-ALLATE 500 EC Selective Herbicide will reduce Wild oat control. Control may be poor in the first years of practicing no-till/min till. To maintain crop safety attention to sowing speed and soil throw is required. <b>Avoid throwing treated soil into adjacent sowing furrows.</b> This is especially critical at higher use rates. <b>Avoid sites that water log or where furrow walls may collapse as crop establishment &amp; vigour may be reduced.</b> Incorporate within 24 hours of application when applied onto a dry soil. Incorporate within 6 hours when applied onto a moist soil. If applied onto a dry soil, and it rains before sowing/incorporation is planned, incorporate within 6 hours to ensure the effectiveness of 4Farmers TRI-ALLATE 500 EC Selective Herbicide is retained.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL  
UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**WITHHOLDING PERIOD-HARVEST:**  
NOT REQUIRED WHEN USED AS DIRECTED.

**WITHHOLDING PERIOD-GRAZING:**

Cereal forage: DO NOT graze or cut for stockfood for 11 weeks after application.  
Oilseed forage: DO NOT graze or cut for stockfood for 13 weeks after application.  
Pulse forage: DO NOT graze or cut for stockfood for 13 weeks after application.

## GENERAL INSTRUCTIONS

### WEED CONTROL

4Farmers TRI-ALLATE 500 EC Selective Herbicide normally provides 6 to 8 weeks control of wild oats when the seedbed is moist during the emergence of wild oats. Under very dry soil surface (top 2 cm) conditions, wild oats may not be controlled. Wild oats may emerge between application and complete incorporation by sowing particularly if initial incorporation is provided by harrows. 4Farmers TRI-ALLATE 500 EC Selective Herbicide may not control wild oats which have germinated at the time of application and incorporation if the top 2 cm of soil is dry. 4Farmers TRI-ALLATE 500 EC Selective Herbicide will not control emerged wild oats.

### CROP SAFETY

Reduced emergence of cereals and linseed may occur if seed is placed in treated soil layer. Do not plant cereal crops at greater than 5 to 7 cm. Overspraying of headlands or corners may result in crop damage. On heavy clay soils, crop thinning may occur in wheel tracks. Environmental conditions which adversely affect emergence or early crop growth, may predispose crops to injury, particularly where tank mixes with other pre-emergence herbicides are used. Read and follow label directions, restraints, plantback periods and safety directions for tank mix products.

### SOIL PERSISTENCE AND FOLLOWING CROPS

Under prolonged and extremely dry conditions 4Farmers TRI-ALLATE 500 EC Selective Herbicide may persist for several months. Damage to field oats and sorghum may occur.

### MIXING

4Farmers TRI-ALLATE 500 EC Selective Herbicide is an oily concentrate that mixes readily with water. Add the required amount of 4Farmers Tri-Allate 500 EC Selective Herbicide to half the required amount of water in the spray tank. Mix thoroughly and add the remaining water.

### APPLICATION

Apply to soil in friable condition so that uniform incorporation of the product can be achieved to a depth of 2 to 4 cm. The soil should be free of large clods (>75 mm) and should not have crop or weed residue at a level which will interfere with the free flow of soil through planting equipment providing full soil disturbance. Emerged wild oats should be controlled either by prior cultivation/application of a knockdown herbicide or by the tank mixing of a knockdown herbicide with 4Farmers TRI-ALLATE 500 EC Selective Herbicide. Apply 4Farmers TRI-ALLATE 500 EC Selective Herbicide by ground sprayer in 30 to 100 litres of water per hectare using a boomspray fitted with flat fan nozzles. Use spraying pressure of approximately 200-240 kPa. Avoid application under strong and/or gusty wind conditions. DO NOT apply by aircraft.

### INCORPORATION

Two pass incorporation provides superior weed control to single pass incorporation on heavier textured soils.

- (i) 4Farmers TRI-ALLATE 500 EC Selective Herbicide may be incorporated initially in a suitable cultivated seedbed with either heavy 5 row harrows e.g. Phoenix harrow, or tined cultivator. Final incorporation with a combine or airseeder/cultivator is essential to provide for maximum weed control. Heavy tined harrows or a rotary harrow should trail the planter or follow as a separate operation at an angle to the planter to complete incorporation. Ensure that the soil surface is left free of pronounced ridging.

- (ii) On soils in friable condition, 4Farmers TRI-ALLATE 500 EC Selective Herbicide may be incorporated with the planting operation only up to 24 hours following application onto a dry soil surface or up to 6 hours on moist surface soil. If 4Farmers TRI-ALLATE 500 EC Selective Herbicide has been sprayed on dry soil, and it rains before incorporation is planned, the product must be incorporated within 6 hours to retain its herbicide effectiveness. Planting should be carried out with a combine or airseeder/cultivator which provides full soil disturbance. Heavy tined harrows or a rotary harrow should trail the planter or follow as a separate operation to complete incorporation. Ensure that the soil surface is left free of pronounced ridging. Do not operate incorporation equipment at greater than 5 to 7 cm depth. Where soil is wet at planting, poor incorporation may result. For shallow-sown legume and oil-seed crops, complete incorporation by cultivation to a depth of 5 cm must be undertaken before planting. Note that point size should be matched to tyne spacing to avoid uneven incorporation. For example, a wideline seeder with 15 cm tyne spacing should be fitted with unworn 15 cm points. A combine with 9 cm spacing should be fitted with an evenly worn set of 10 cm points. Sweeps or other low profile points may not provide satisfactory incorporation.

### COMPATIBILITY

4Farmers TRI-ALLATE 500 EC Selective Herbicide is compatible with: 4FARMERS CHLORSULFURON 750 WDG WATER DISPERSIBLE GRANULES, 4FARMERS DICLOFOP-METHYL 375 SELECTIVE HERBICIDE 4FARMERS DICLOFOP-METHYL 500 EC SELECTIVE HERBICIDE, 4FARMERS TRIFLURALIN 480 SELECTIVE HERBICIDE, 4FARMERS GLYPHOSATE 470 HERBICIDE, 4FARMERS TRIFLURALIN-ORYZALIN 250 EC SELECTIVE HERBICIDE.

### RESISTANT WEEDS WARNING

GROUP	J	HERBICIDE
-------	---	-----------

4Farmers TRI-ALLATE 500 EC Selective Herbicide is a member of the thiocarbamate group of herbicides. 4Farmers TRI-ALLATE 500 EC Selective Herbicide has the inhibition of mitosis mode of action. For weed resistance management 4Farmers TRI-ALLATE 500 EC Selective Herbicide is a Group J herbicide. Some naturally occurring weed biotypes resistant to 4Farmers TRI-ALLATE 500 EC Selective Herbicide and other Group J herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These weeds will not be controlled by 4Farmers TRI-ALLATE 500 EC Selective Herbicide or other Group J herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, 4Farmers Australia Pty Ltd accepts no liability for any losses that may result from the failure of 4Farmers TRI-ALLATE 500 EC Selective Herbicide to control resistant weeds.

### PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

### PROTECTION OF LIVESTOCK, FISH, CRUSTACEA AND ENVIRONMENT

Dangerous to fish. Do not contaminate dams, rivers or streams with 4Farmers TRI-ALLATE 500 EC Selective Herbicide or used containers.

### STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

### Refillable Containers (110L and 1000L)

Empty contents fully into application equipment and return to point of sale for refill or storage.

### SAFETY DIRECTIONS

May irritate eyes and skin. Avoid contact with eyes, skin and clothing. Do not inhale spray mist. When preparing spray, wear cotton overalls buttoned to the neck and wrist, washable hat, elbow-length PVC gloves and face shield or goggles. If product on skin immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms, and face thoroughly with soap and water. After each day's use wash gloves, face shield and contaminated clothing.

### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. If swallowed, DO NOT induce vomiting. Give a glass of water.

### SAFETY DATA SHEET

For further information please refer to the Safety Data Sheet. For a copy visit our website at [www.4farmers.com.au](http://www.4farmers.com.au)

### CONDITIONS OF SALE

"4Farmers Australia Pty Ltd ('4Farmers') shall not be liable for any loss injury damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on 4Farmer's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of 4Farmers has any authority to add to or alter these conditions."

<sup>1</sup> Not a Trademark of 4Farmers Australia Pty Ltd

In a Transport Emergency  
Dial 000  
Police or Fire Brigade

Special Poisons Advice:  
13 11 26