

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



EPOXICONAZOLE 125 FUNGICIDE

ACTIVE CONSTITUENT: 125 g/L EPOXICONAZOLE

GROUP 3 FUNGICIDE

For the control of certain fungal diseases including stripe rust of wheat and specified fungal diseases of barley, as per the Directions for Use table.

**IMPORTANT: READ THIS LEAFLET THOROUGHLY
BEFORE USING THIS PRODUCT**

4 FARMERS AUSTRALIA PTY LTD

A.B.N. 51 160 092 428

1 / 70 MCDOWELL STREET, WELSHPOOL, WA 6106

TEL: (08) 9356 3445 www.4farmers.com.au

Batch Number:

Date of Manufacture:

APVMA Approval No: 65949/52374

DIRECTIONS FOR USE

CROP	DISEASE	RATE PER HA	CRITICAL COMMENTS
Wheat	Leaf rust (<i>Puccinia recondita</i>)	500 mL	Apply when conditions favour disease development and prior to development of disease in the crop. Aim to apply between stem elongation and ear emergence complete (ZGS 32-59) if disease is present. DO NOT apply later than ZGS 59.
	Stripe rust (<i>Puccinia striiformis</i>) Septoria nodorum blotch (<i>Phaeosphaeria nodorum</i>)	250 to 500 mL	Apply when conditions favour disease development and prior to development of disease in the crop. Aim to apply between stem elongation and ear emergence complete (ZGS 32-59) if disease is present. DO NOT apply later than ZGS 59. Apply the higher rate when disease is present on the top leaf or conditions are favourable for disease development at these stages of crop development. Repeat spraying may be required, particularly if infection occurs early. Regularly monitor the crop from 3-4 weeks after the first application for signs of re-infection.
	Powdery mildew (<i>Blumeria graminis f.sp. tritici</i>)	250 mL	Apply when conditions favour disease development and prior to development of disease in the crop. Aim to apply during early tillering (ZGS 21-22) stage of the crop if disease is present. DO NOT apply later than ZGS 59. Repeat spraying may be required, particularly if infection pressure persists. Regularly monitor the crop from 3-4 weeks after the first application for signs of re-infection.
Barley			DO NOT exceed a total rate of 500 mL/ha of 4FARMERS EPOXICONAZOLE 125 in a growing season.
	Leaf rust (<i>Puccinia hordei</i>)	250 to 500 mL	Apply when conditions favour disease development and prior to development of disease in the crop. Aim to apply between stem elongation and ear emergence complete (ZGS 32-59) if disease is present. DO NOT apply later than ZGS 59. Apply the higher rate when disease is present on the top leaf or conditions are favourable for disease development at these stages of crop development.
	Net form of net blotch (<i>Pyrenophora teres f. sp. teres</i>)	250 to 500 mL	Apply when conditions favour disease development and prior to development of disease in the crop. Aim to apply between stem elongation and ear emergence complete (ZGS 32-59) if disease is present. DO NOT apply later than ZGS 59. Two applications of 4FARMERS EPOXICONAZOLE 125 at the lower rate may provide better control than a single application under certain conditions. If applying a two spray program, only apply 250 mL/ha per application, with the first application at ZGS 31-32 if the disease is present. Aim to apply the second application around ZGS 37-39, but no later than ZGS 59, if infection pressure persists. DO NOT exceed a total rate of 500 mL/ha of 4FARMERS EPOXICONAZOLE 125 in barley per growing season.
	Leaf scald (<i>Rynchosporium secalis</i>) Powdery mildew (<i>Blumeria graminis f. sp. hordei</i>)	250 mL	Apply when conditions favour disease development and prior to development of disease in the crop. Aim to apply between early tillering and flag leaf emergence (ZGS 21-39) if disease is present. DO NOT apply later than ZGS 59. Repeat spraying may be required, particularly if infection pressure persists. Regularly monitor the crop from 3-4 weeks after the first application for signs of re-infection. DO NOT exceed a total rate of 500 mL/ha of 4FARMERS EPOXICONAZOLE 125 in barley per growing season.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:

DO NOT HARVEST FOR 6 WEEKS AFTER APPLICATION.

DO NOT GRAZE OR CUT FOR STOCKFOOD FOR 6 WEEKS AFTER APPLICATION.

GENERAL INSTRUCTIONS

4FARMERS EPOXICONAZOLE 125 is a broad spectrum contact and systemic fungicide. It is taken up via the stem and foliage and is translocated upwards and outwards to provide protection to new growth. Thorough coverage of the crop is necessary for best results. Check the crop regularly for any symptoms of disease infection. Apply as soon as disease is present as indicated in the Directions for Use.

MIXING

4FARMERS EPOXICONAZOLE 125 is a suspension concentrate formulation. The addition of a non-ionic surfactant is generally not required, although under certain environmental conditions and lower total application volumes, the addition of Chemwet 1000 or other recommended surfactant may assist in providing better coverage of sprayed surfaces. Add the product to the half filled spray tank while agitating. Continue to agitate while topping up the tank and during spraying.

APPLICATION

Ground Application

Apply in a water volume of between 50 and 100L/ha, using flat fan nozzles operating at around 50cm above the top of the crop. Use the higher water volume in crops with heavier canopies.

Aerial Application

Apply with suitable aircraft, set up and operated to apply fungicides to cereal crops in a minimum water volume of 20L/ha.

COMPATIBILITY

For information on compatibility please contact your local reseller or 4Farmers Australia Pty Ltd representative.

FUNGICIDE RESISTANCE WARNING

GROUP 3 FUNGICIDE

4FARMERS EPOXICONAZOLE 125 is a member of the DMI group of fungicides. For fungicide resistance management the product is a Group 3 fungicide. Some naturally occurring individual fungi resistant to the product and other Group 3 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by this product or other Group 3 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, 4Farmers Australia Pty Ltd accepts no liability for any losses that may result from the failure of this product to control resistant fungi.

PRECAUTIONS

Re-entry period: DO NOT allow entry into treated areas until the spray has dried.

CAUTION

Export of treated cereals

Growers should note that Maximum Residue Limits (MRLs) or import tolerances do not exist in all markets for cereals treated with 4FARMERS EPOXICONAZOLE 125 Fungicide. Additionally, some export markets have established MRLs different to those in Australia. If you are growing cereals for export, please check with 4Farmers Australia Pty Ltd or your grain exporter for the latest information on MRLs and import tolerances BEFORE using 4FARMERS EPOXICONAZOLE 125 Fungicide.

LIVESTOCK DESTINED FOR EXPORT MARKETS

The grazing withholding period only applies to stock slaughtered for the domestic market. Some export markets apply different standards. To meet these standards, ensure that in addition to complying with the grazing withholding period, the Export Slaughter Interval is observed before stock are sold or slaughtered.

EXPORT SLAUGHTER INTERVAL (ESI):

AFTER OBSERVING THE WITHHOLDING PERIOD FOR GRAZING OR CUTTING FOR STOCKFOOD, LIVESTOCK THAT HAVE BEEN GRAZED ON OR FED TREATED CROPS SHOULD BE PLACED ON CLEAN FEED FOR 3 DAYS PRIOR TO SLAUGHTER.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET CROPS

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Highly toxic to aquatic organisms. DO NOT apply under weather conditions or from spraying equipment which could be expected to cause spray drift onto adjacent crops, crop lands, livestock, pastures, wetlands, waterbodies or watercourse.

DO NOT apply 4FARMERS EPOXICONAZOLE 125 aerially if waterbodies, watercourses or wetlands are within 30 metres downwind of the application area. Do NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

SAFETY DIRECTIONS

Harmful if inhaled or swallowed. Will irritate the eyes, nose, throat and skin. Avoid contact with eyes and skin. Do not inhale vapour. When opening the container and preparing the spray wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow-length PVC gloves and a half facepiece respirator. When using the prepared spray wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and a washable hat. If product on skin, immediately wash area with soap and water. Wash hands after use. After each day's use, wash gloves, contaminated clothing and respirator and if rubber wash with detergent and warm water.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

MATERIAL SAFETY DATA SHEET

For further information please refer to the Safety Data Sheet. For a copy visit our website at www.4farmers.com.au

CONDITIONS OF SALE

"Any provisions or rights under the Trade Practices Act 1974 or relevant state legislation which cannot be excluded by those statutes or by law are not intended to be excluded by these conditions of sale. Subject to the foregoing, all warranties, conditions, rights and remedies, expressed or implied under common law, statute or otherwise, in relation to the sale, supply, use or application of this product, are excluded. 4Farmers Australia Pty Ltd and/or its affiliates ("4Farmers") shall not accept any liability whatsoever (including consequential loss), or howsoever arising (including negligence) for any damage, injury or death connected with the sale, supply, use or application of this product except for liability which cannot be excluded by statute."

In a Transport Emergency Dial 000 Police or Fire Brigade	Special Poisons Advice: 13 11 26
---	---