

DANGEROUS POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



FENAMIPHOS 400

Nematicide and Insecticide

ACTIVE CONSTITUENT: 400 g/L FENAMIPHOS
(An Anticholinesterase Compound)

GROUP 1B INSECTICIDE

For control of nematodes on aloe vera planting material
and banana planting material as specified in the
Directions for Use Table.

**READ THIS INFORMATION THOROUGHLY BEFORE
USING THIS PRODUCT**

4 FARMERS AUSTRALIA PTY LTD

A.B.N. 51 160 092 428

1 / 70 McDowell Street, Welshpool, WA 6106

Tel: (08) 9356 3445 Fax: (08) 9356 3447

Batch Number:

Date of Manufacture:

APVMA Approval No: 62574/0715

RESTRAINTS

DO NOT dispose of spent dip on site. Spent dip waste must be transported to a facility that can lawfully receive it.

DIRECTIONS FOR USE

Directions for use table attached at end of document.

OTHER LIMITATIONS

WOOD DUCKS (*Chenonetta jubata*) MAY BE PUT AT RISK IF THIS PRODUCT IS USED INCORRECTLY.

DO NOT USE THIS PRODUCT IN THE HOME GARDEN.

WITHHOLDING PERIODS

BANANA AND ALOE VERA PLANTING MATERIAL: NOT REQUIRED WHEN USED AS DIRECTED.

TRADE ADVICE

EXPORT OF TREATED PRODUCE

Growers should note that MRLs or import tolerances **DO NOT** exist in all markets for edible produce treated with 4Farmers Fenamiphos 400 Nematicide and Insecticide. If you are growing edible produce for export, please check with the relevant regulatory authorities for the latest information on MRLs and import tolerances before using 4Farmers Fenamiphos 400 Nematicide and Insecticide.

GENERAL INSTRUCTIONS

MIXING

Add the required quantity of 4Farmers Fenamiphos 400 Nematicide to the dip vat full of water while stirring. Agitate before use if mixture has been left overnight.

ACCELERATED BIODEGRADATION

As with other soil insecticides the effectiveness of this product may be reduced as a result of the soil microflora developing the ability to biodegrade at an accelerated rate, particularly after repeated application.

If unacceptable control occurs, immediately contact your 4Farmers Australia Pty Ltd representative for detailed information. This product should not be used where the user suspects the soil microflora have developed the ability to degrade it at an accelerated rate.

4Farmers Australia Pty Ltd accepts no responsibility, and intends by this statement to exclude liability, if the effectiveness of this product is reduced or nullified as a result of accelerated biodegradation, since biodegradation is accelerated by soil characteristics which occur naturally and are beyond 4Farmers Australia Pty Ltd's control.

RESISTANCE WARNING

GROUP 1B INSECTICIDE

For insecticide resistance management 4Farmers Fenamiphos 400 Nematicide and Insecticide is a Group 1B insecticide.

Some naturally occurring insect biotypes resistant to 4Farmers Fenamiphos 400 Nematicide and Insecticide may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if 4Farmers Fenamiphos 400 Nematicide and Insecticide and other Group 1B insecticides are used repeatedly. The effectiveness of 4Farmers Fenamiphos 400 Nematicide and Insecticide on resistant individuals could be significantly reduced. Since the occurrence

of resistant individuals is difficult to detect prior to use, 4Farmers Australia Pty Ltd accepts no liability for any losses that may result from the failure of 4Farmers Fenamiphos 400 Nematicide and Insecticide to control resistant insects.

4Farmers Fenamiphos 400 Nematicide and Insecticide may be subject to specific resistance management strategies. For further information contact your local suppliers, 4Farmers Australia Pty Ltd representative or local agricultural department agronomist.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Dangerous to fish and other aquatic vertebrates. **DO NOT** contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in a locked room or place away from children, animals, food, feedstuffs, seed and fertilisers. Store in the closed, original container in a dry, cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight. Triple-rinse containers before disposal. Add rinsings to dip tank. **DO NOT** dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. **DO NOT** burn empty containers or product.

DIP DISPOSAL

DO NOT dispose of spent dip on site. Spent dip waste must be transported to a facility that can lawfully receive it.

SAFETY DIRECTIONS

Poisonous if absorbed by skin contact or inhaled or swallowed.

Repeated minor exposure may have a cumulative poisoning effect.

Will damage eyes and skin. Will irritate the nose and throat.

Avoid contact with eyes and skin. Do not inhale vapour.

When preparing dip and using the prepared dip wear cotton overalls buttoned to the neck and wrist and a washable hat, protective waterproof clothing or a PVC apron, elbow-length chemical resistant gloves and impervious footwear and face shield or goggles.

If clothing becomes contaminated with product or wet with spray remove clothing immediately.

If product in eyes, wash it out immediately with water.

If product or dip on skin, immediately wash area with soap and water.

After using and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water.

After each day's use, wash gloves and goggles and contaminated clothing.

FIRST AID INSTRUCTIONS

If swallowed, splashed on skin or in eyes, or inhaled, contact the Poisons Information Centre (phone Australia 13 11 26) or a doctor at once.

Remove any contaminated clothing and wash skin thoroughly.

If swallowed, activated charcoal may be advised. Give atropine if instructed.

DIRECTIONS FOR USE:

| CROP | PEST | STATE | RATE | CRITICAL COMMENTS |
|-----------------------------|--------------------------------------|-------------------|-------------------|--|
| Aloe Vera planting material | Soil borne plant parasitic nematodes | QLD, NSW, WA only | 1 L/100L water | Air dry or cure prepared pups for 48 hours prior to treatment. Immerse unpared pups in 4Farmers Fenamiphos 400 Nematicide and Insecticide dip solution for 30 minutes. Allow to drain and dry. Wear elbow-length chemical resistant gloves when dipping and when handling dipped material. |
| Banana planting material | Soil borne plant parasitic nematodes | QLD, NSW, WA only | 100 mL/100L water | Pare the planting material to remove all dead tissue, immerse planting material in 4Farmers Fenamiphos 400 Nematicide and Insecticide dip solution for 10 minutes. Allow to drain and dry. Wear elbow-length chemical resistant gloves when dipping and when handling dipped material. |

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER,
CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER
APPROPRIATE LEGISLATION**

**In a Transport Emergency
Dial 000
Police or Fire Brigade**

**Special Poisons Advice:
13 11 26**