

- **Metsulfuron-methyl** For knockdown weed control in fallows and prior to planting certain winter cereals. Observe crop safety/crop rotation recommendations.
- **Oxyfluorfen** (Goal CT®) Mixture requires a specific compatibility agent, seek advice from labels.
- **Note:** Addition of crystalline (not prilled or granular) ammonium sulphate at the rate of 2kg/100L spray solution is required to overcome antagonism. This mixture is also useful for improving the performance of 4Farmers Glyphosate 450 under adverse environmental conditions such as cool cloudy weather. Ammonium sulphate may be corrosive to metal parts of the sprayer.

Insecticides:

- **Dimethoate, Omethoate, Fenitrothion, Phosmet (Imidan®), Chlorpyrifos**
- Other insecticides have not been compatibility tested

APPLICATION INFORMATION

- 4Farmers Glyphosate Herbicide is a non-selective, translocated herbicide.
- Direct contact, or even drift, may cause severe injury or destruction to any growing crop or other desirable plants including trees.

Boom equipment:

- Application of this product in low spray volumes (25-100L/ha) is recommended.
- Fan nozzle equipment is recommended, using pressures in the range 240-280kPa.
- Boom height must be set to ensure double overlap of nozzle patterns at the top of the weed canopy.

Aerial Equipment:

- Aerial application should only be in pasture or fallow crop situations prior to establishment of field or fodder crops or new pasture and for pre harvest application for sorghum crops.
- **DO NOT** use in intensive cropping situations.
- Use registered rates up to a maximum of 3.2L/ha and for Micronair and boom equipment apply in a minimum spray volume of at least 15L/ha.
- Average droplet size should be 250-350 micron diameter with a swath width of 15-17 metres.
- **DO NOT** apply by aircraft in temperatures above 35°C.
- Thoroughly wash aircraft, especially landing gear, after each day of spraying, to remove herbicide.

Application on hilly terrain:

- Increase water volume to 30-80L/ha with droplet size to at least 300 micron VMD on hilly terrain.

Application under hot conditions:

- When applying in temperatures above 25°C, increase water volume to at least 30L/ha and droplet size to at least 300 micron VMD.
- Avoid conditions which may result in drift such as wind over 8km/h and inversion conditions or equipment producing droplets of 150 micron or less.

MAINTENANCE AND CLEANING

- Thoroughly flush tanks, pumps and nozzles with clean fresh water after each day's spraying.

PRECAUTION

- **DO NOT** mix, store or apply this product or spray solutions of this product in galvanised or unlined steel containers or spray tanks, since a highly flammable gas may be formed. Use stainless steel, aluminium, brass, copper, fibreglass, plastic or plastic-lined containers or spray tanks.

PROTECTION OF CROPS, NATIVE AND OTHER NON-

TARGET PLANTS

- **DO NOT** apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.
- **DO NOT** apply prior to transplanting tomato seedlings.

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND

ENVIRONMENT

- **DO NOT** contaminate dams, rivers or streams with the product or used container.
- **DO NOT** apply to weeds growing in or over water. **DO NOT** spray across open bodies of water.

PROTECTION OF LIVESTOCK

- **DO NOT** graze or cultivate for one day after application (annual weeds) or 7 days (perennial weeds) to ensure herbicide absorption.
- There is no withholding period for grazing stock. However, it is recommended that grazing of treated plants be delayed as above. Certain plants, eg, Soursob, St John's wort, Bracken, may be toxic to stock. Grazing should be delayed until complete browning of these plants has occurred.

STORAGE AND DISPOSAL (20L)

- Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.
- Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

STORAGE AND DISPOSAL (110L)

- Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.
- Store this tightly closed minibulk container in a safe and secure place away from children, animals, food and feedstuffs. Storage must be secure so that contents cannot be tampered with. All locks and/or seals must be in order. If locks or seals are broken prior to initial use then the integrity of this product cannot be assured. If this occurs 4 Farmers Australia Pty Ltd should be advised immediately. This minibulk container is reusable and remains the property of 4 Farmers Australia Pty Ltd.
- **DO NOT** rinse empty container. Empty contents fully into application equipment. Close all valves and return to the point of supply for refill or storage. No other liquid, solid or pesticide product should be put into it. When empty return to 4Farmers Pty Ltd for cleaning, relabelling and refilling.

SAFETY DIRECTIONS

- Product will irritate eyes and skin.
- Avoid contact with eyes and skin.
- When preparing product for use, wear elbow-length PVC gloves and face shield (or goggles).
- When using controlled droplet applicator, wear protective waterproof clothing and impervious footwear.
- After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water.
- After each day's use wash contaminated clothing, gloves and face shield or goggles.

FIRST AID

- If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.

SAFETY DATA SHEET

For further information please refer to the Safety Data Sheet which can be obtained from www.4farmers.com.au

CONDITIONS OF SALE

The use of 4Farmers Glyphosate 450 being beyond the control of the manufacturer, no warranty expressed or implied is given by 4 Farmers Australia Pty Ltd regarding its suitability, fitness or efficacy for any purpose for which it is used by the buyer, whether in accordance with the directions or not and 4 Farmers Australia Pty Ltd accepts no responsibility for any consequences whatsoever resulting from the use of this product.

NOT CLASSIFIED AS DANGEROUS GOODS FOR TRANSPORT BY ROAD AND RAIL

¹ Registered trademark of Velsicol Chemical Corp., USA

² Registered trademark of Rohm and Haas Co., USA

³ Registered trademark Bayer, Germany

⁴ Registered trademark of Stauffer Chemical Company, USA

In a Transport Emergency Dial 000 Police or Fire Brigade	Special Poisons Advice: 13 11 26
---	---