

## GENERAL INSTRUCTIONS

In **Wheat** Triadimenol+ Seed Dressing, especially at the higher rate of 150 gm/100 Kg seed, may delay emergence. Crop will recover rapidly in moist soil types. However, seed treated with the 150 gm/100 Kg seed, should not be sown in soils that are likely to clod or crust over after sowing. Sowing Triadimenol+ treated wheat deeper than 50 mm may slightly reduce percentage plant establishment in heavier soils and sowing rates may need to be adjusted accordingly. The possible delay in emergence of wheat treated with the 150 gm/100 Kg seed rate may counteract the advantage of suppression of speckled leaf blotch.

Treated seed may be held over for sowing in the following season. If so, additional insecticide treatment may be necessary. Treated seed not used for sowing should be destroyed. If the seed is not used immediately after treatment, it should be stored in a dry, well-ventilated place.

## FUNGICIDE RESISTANCE WARNING

GROUP **C** FUNGICIDE

Triadimenol is a member of the D.M.I. group of fungicides. For fungicide resistance management, 4Farmers Triadimenol 150 is a Group C fungicide.

Some naturally occurring individual fungi resistant to 4Farmers Triadimenol 150 and other Group C fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by 4Farmers Triadimenol 150 and other Group C fungicides, thus resulting in a reduction in efficacy and possible yield loss.

Since the occurrence of resistant fungi is difficult to detect prior to use, 4Farmers Pty Ltd accepts no liability for any losses that may result from the failure of 4Farmers Triadimenol 150 to control resistant fungi.

### MIXING as a powder

The required quantity of Triadimenol+ seed dressing should be thoroughly mixed with the correct weight of seed in a seed dressing machine or a suitable revolving drum, barrel or auger. It is recommended to wear a face mask while handling the dust and treated seed.

## DIRECTIONS FOR USE

Restraint: DO NOT treat physically damaged seed.

Crop	Disease/Pest	State	Rate	Critical Comments
Wheat	Bunt, Flag Smut (seed and soil borne), Loose Smut. Suppression of seedling infections of Stripe Rust.	Old, NSW, Vic, Tas, SA WA only	100g/100kg seed	Thorough mixing is essential. Ensure that the recommended rate is applied.
Oats,	Covered and Loose Smut.			
Barley	Covered and Loose Smut, suppression of seedling infections of Powdery Mildew. Suppression of seedling infections of Leaf Scald.	NSW, Vic, Tas, SA, WA only		
Wheat	Bunt, Flag Smut (seed and soil borne), Loose Smut. Suppression of seedling infections of Stripe Rust.	NSW, Vic, Tas, SA, WA only	150g/100kg seed	Use the 150g/100kg seed rate in areas where foliar diseases frequently occur and where any of the following are required: – Longer suppression of stripe rust on seedlings – Control of seedling infections of powdery mildew – Suppression of speckled leaf blotch wheat – Extended suppression of leaf scald in barley: and use higher rate on more susceptible varieties or where disease pressure is heavy.
	Suppression of seedling infections of Speckled Leaf Blotch.	NSW, Vic, Tas, SA only		
Barley	Covered and Loose Smut. Suppression of seedling infections of Powdery Mildew. Suppression of seedling infections of Leaf Scald.	NSW, Vic, Tas, SA only		
Wheat, Barley, Oats	Protection against insect pests of stored grains: Granary Weevil, Indian Meal Moth, Lesser Grain Borer, Rice Weevil, Rust-Red Flour Beetle, Sawtooth Grain Beetle, Tropical Warehouse Moth.	Old, NSW, Vic, Tas, SA, WA only	100 or 150g/100kg seed	Both rates give protection against insect pests. Use rate appropriate to disease control required.

**NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**WITHHOLDING PERIOD: DO NOT GRAZE OR CUT FOR STOCK FEED WITHIN 5 WEEKS OF SOWING.**

## MIXING for use in a pickle pot or similar (liquid application)

- The required quantity of 4Farmers Triadimenol 150 W.P. should be thoroughly mixed with a small quantity of clean, filtered water to make a creamy mix. Add this blend to the required quantity of water in the tank and mix thoroughly. Agitate prior to and during spraying.
- DO NOT mix more than can be used for the day's seeding.
- DO NOT tank mix with trace elements or other chemicals without first checking compatibility.

## PROTECTION OF LIVESTOCK

- Seed treated with this product must not be used for animal consumption, poultry feed or mixed with animal feed.
- DO NOT mix seed treated with this product with grain intended for animal consumption.

## PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND THE ENVIRONMENT

- DO NOT feed treated leaf matter to wild or domestic birds or other animals.
- Harmful to fish.
- DO NOT contaminate streams, rivers or waterways with the chemical or used containers.
- Used containers or bags which have held treated seed must be correctly disposed of.

## Storage and Disposal and Protection of Others

When treated seed is stored it should be kept apart from other grain. Such bags or other containers should be clearly marked to indicate the contents have been treated with Triadimenol+.

Store the unused product in the closed original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

Do not allow seed treated with this product to contaminate grain intended for human consumption.

Do not use seed treated with Triadimenol+ for human consumption.

Bags which have held treated seed with this product are not to be used for any other purpose. Shake empty bag. If using as a liquid, single rinse the bag before disposal and add rinsings to the spray tank. Do not dispose of undiluted chemicals on-site. Puncture and bury empty containers

in a local authority land fill. If no land fill available, bury the containers to a depth of 500 mm in a disposal area specifically marked and set up for this purpose clear of waterways desirable vegetation and tree roots. Empty containers and product should not be burnt.

## SAFETY DIRECTIONS

- **Poisonous if swallowed.**
- May irritate the eyes and skin.
- Facial skin contact may cause temporary facial numbness.
- Avoid contact with eyes and skin.
- Do not inhale dust. Wear a face mask while handling the dust and treated seed.
- Wash hands, arms and face thoroughly with soap and water after use and before eating, drinking or smoking.

## FIRST AID

- If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.
- If swallowed, and if more than 15 minutes from a hospital, induce vomiting, preferably using Ipecac Syrup APF.
- For further information see Material Safety Data Sheet which can be obtained from the supplier.

## Limitation of Liability

This product was designed only for the particular purpose indicated on the label. Failure to use this product strictly as directed may be illegal; prove dangerous; render the product ineffective and avoid any Warranty or Liability of 4Farmers Pty Ltd.

## CONDITIONS OF SALE

The use of 4Farmers Triadimenol 150 W.P., being beyond the control of the manufacturer, no warranty expressed or implied is given by 4Farmers Pty. Ltd. regarding its suitability, fitness or efficacy for any purpose for which it is used by the buyer, whether in accordance with the directions or not and 4Farmers Pty. Ltd. accepts no responsibility for any consequences whatsoever resulting from the use of this product.

**IN A TRANSPORT EMERGENCY  
DIAL 000  
POLICE OR THE FIRE BRIGADE**

NRA Approval Number: 49137/0498

Batch Number:

Date of Manufacture: