

Bromacil 800

Similar to: *Uragan*[®]

Active Constituent	Bromacil 800g/kg		Formulation	WG
Typical Situations	Horticultural crops, around agricultural buildings, industrial, commercial, rights of way and other (see registration section).			
Chemical Group	C	Mode of Action	Root and some foliar uptake, upward translocation – disrupts photosynthesis, destroys cells.	
Price ex GST	\$40.00/kg	Typical Pack Size	5kg	
Poison Schedule	N/S*	Dangerous Goods Class	-	UN -
Withholding Period	Harvest	N/A		
	Grazing	N/A		
Plant Back	3 years, everything except pineapples.			
Application Method	Boom spray 50L/ha minimum, best at 250-450L/ha			Rain fastness
				N/A
Efficacy	Sufficient moisture to carry the chemical into the soil is required within 3-4 days for best efficacy.			
Adjuvants	Wetter 1000 (if weeds present).			
Compatibility	Most herbicides (confirm individual mixes using a jar test).			
Incompatibility	Grass selectives, Group B herbicides, Simazine.			
Water Quality	N/A			
Time to Effects and Symptoms	7-10 days - chlorosis and desiccation in sunny conditions.			
4F Broadacre Registrations	N/A			
4F Other Registrations	Citrus, asparagus, pineapple, industrial and rights of way and other (extract listed).			
Similar Product Registrations	<i>Uragan</i> [®] is registered for fence lines.			
Situation	Target Weed/s	Rate/ha	Comments	
Citrus (established groves only)	Most annual weeds	2.2-4.5kg	Vary rate with soil type (e.g. light soil 2.2kg, heavy soil 4.5kg). Apply to a weed free ground in Autumn to Spring.	
	Perennial grasses, e.g. Couch grass, Nut grass, Johnson grass, Kikuyu, Skeleton weed	4.5-6.5kg		
Industrial Areas (around sheds, etc.)	Barley grass, Bathurst Burr, Brome grass, Caltrop, Capeweed, Cobblers peg, Doublegee, Paterson's Curse, Prairie grass, Annual Ragweed, Annual Ryegrass, Saffron thistle, Summer grass, Thornapple, Wild Mustard, Wild oats, Wild Radish, Wild Turnip, Winter grass and other (see label)	3.5-6.5kg	Lower rates are for lighter soils, higher rates for heavier soils. Rates will also increase with increase in rainfall with the lowest rate for 250mm or less per year. Apply retreatment when grasses reappear. Use Wetter 1000 where weeds are present.	
	Bahia grass, Bluetop, Guinea grass, Horehound, Para grass, Purple Top, Plantain, St John's Wort, Soursob	8-13kg		
	Bindweed, Couch grass, Dock, Johnson grass, Kikuyu, Nut grass, Skeleton weed, Sorrel, Water Couch	16.5-22kg		
	Retreatment for above	2-6.5kg		
Comments				
Do not apply more than 9kg/ha per year in citrus and pineapple crops. In low rainfall environments deep rooted perennials may not be controlled. Commonly used in combination with Paraquat for Glyphosate-resistant weed control on firebreaks and other non-agricultural areas and provides long periods of soil residual activity. *N/S – Not scheduled				