



Product Name: 4FARMERS TAKE UP
APVMA Approval No: 87210 / 119226

Label Name:	4FARMERS TAKE UP
-------------	------------------

Signal Headings:	CAUTION KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING
------------------	--

Constituent Statements:	637g/L Liquid Hydrocarbon 240g/L Non-IONIC Surfactants
-------------------------	---

Mode of Action:	
-----------------	--

Statement of Claims:	AN ADJUVATED SPRAYING OIL TO IMPROVE THE WETTING AND SPREADING OF HERBICIDES, INSECTICIDES AND DEFOLIANTS ON PLANT SURFACES AS PER DIRECTION FOR USE TABLE.
----------------------	---

Net Contents:	5L to 1000L
---------------	-------------

Restrains:	
------------	--

Directions for Use:	This section contains file attachment.
---------------------	--

Other Limitations:	
--------------------	--

Withholding Periods:	Adhere to the withholding period(s) specified on the Label of the companion product(s) used.
----------------------	--

Trade Advice:	
---------------	--

General Instructions:	<p>MIXING: • These are general mixing instructions. Always follow the mixing instructions of each ingredient in the tank mix. Ensure product compatibility. Recommendations on other product labels always take precedence over mixing instructions on this label. • Add water to the spray tank to 10cm above the level of agitation and ensure the agitation device is working vigorously. (There must be a minimum of 100L of water in the tank before any pesticide is added.) • If tank mixing, use the following general order for mixing partners. First add water dispersible granules (eg Eclipse), second add flowable formulations (eg atrazine) , third add emulsifiable concentrates (eg Fluroxypyr 200EC), fourth add water soluble liquids (eg Clopyralid 300 Herbicide) and last 4Farmers TAKE UP and other adjuvants . • Between the addition of each ingredient allow at least one minute to ensure thorough mixing before the new tank-mix partner is added. • Finally add 4Farmers TAKE UP. Continue filling the tank to the required volume maintaining agitation at all times. • Only mix sufficient spray solution for immediate user. Haloxypop and other tank-mixes should be applied immediately for best results .</p> <p>STORAGE OF SPRAY SOLUTION: Agitate for 1 hour every 8 hours, and agitate for 1 hour before spraying out where spray solution is not applied within 8 hours of mixing.</p> <p>CLEANING SPRAY EQUIPMENT: The presence of the product(s) makes it essential for the spray equipment to be thoroughly cleaned between different applications . To rinse and decontaminate spray equipment refer to the relevant product labels for instructions.</p>
-----------------------	---

Resistance Warning:	
---------------------	--

Precautions:	
--------------	--

Protections:	<p>PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT Do not contaminate streams, rivers or waterways with the chemical or used container.</p>
--------------	---

Storage and Disposal:	<p>STORAGE AND DISPOSAL : Store in the closed original container, in a cool well-ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT store near food, feed stuffs, fertilizers or seed.</p> <p>Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.</p> <p>If not recycling, break, crush, or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.</p> <p>EXPLOSION WARNING : EMPTY containers may contain liquid and/or vapour residue which can be explosive if exposed to ignition source at temperature above 90 degrees Celsius. Such conditions may occur during cutting or welding . DO NOT CUT OR WELD these containers. Explosion may cause injury or death.</p> <p>SMALL SPILL MANAGEMENT: Wear appropriate clothing while cleaning up small spills. Apply absorbent material such as earth, cat litter or clay granules to the spill. Sweep up material for disposal when absorption is complete. Dispose of the contaminated material in accordance with STATE and/or LOCAL regulations .</p>
-----------------------	--

Safety Directions:	<p>SAFETY DIRECTIONS: Will irritate the eyes. May irritate the skin. Avoid contact with eyes and skin. When opening the container and preparing spray wear elbow-length PVC gloves and face shield or goggles. If product in eyes, wash out immediately with water. Wash</p>
--------------------	--

	hands after use. After each day's use wash gloves and face shield. Consult the label of the herbicide(s) used for instructions on protective clothing requirements
--	--

First Aid Instructions:	FIRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766. If swallowed, do NOT induce vomiting.
-------------------------	---

First Aid Warnings:	
---------------------	--

DIRECTIONS FOR USE:

Important : Prior to use read all directions on this label and on the label of the companion crop protection product . The product(s) label for specific usage and restrictions have precedence over directions for use referred to on this label.

4FARMERS TAKE UP is compatible with a wide range of herbicides including: Atrazine#, Eclipse Herbicide, Grazon DS Herbicide, Fluroxypyr 200 Herbicide, Tordon 75-D Herbicide , Clethodim Herbicide, Haloxypop, Broadstrike Herbicide, Triclopyr 600 Herbicide, Clopyralid 750SG Herbicide, Clodinafop 240EC Selective Herbicide , Quizalofop -P- Ethyl Herbicide , Tralkoxydim 400 WG Herbicide. 4FARMERS TAKE UP is compatible with cotton defoliant products such as: Thidiazuron, Ethephon. 4FARMERS TAKE UP is compatible with the insecticides Chlorpyrifos 300.

When mixing with atrazine water dispersible granule formulations, agitation is required.

SITUATION	CRITICAL COMMENTS
Broadacre cropping including fallow (except sugarcane)	Where specified use 500mL 4FARMERS TAKE UP / 100L, applying a minimum of 250mL of 4FARMERS TAKE UP per hectare.
Sugarcane and cotton defoliation	Where specified use 1L 4FARMERS TAKE UP /100L water applying a minimum of 500mL of 4FARMERS TAKE UP per hectare .
Spot and High Volume Spraying	Where specified use 500ml 4FARMERS TAKE UP /100L water applying to point of run-off.
Horticulture	4FARMERS TAKE UP with Haloxypop Herbicide for horticultural uses. Use 500mL 4FARMERS TAKE UP /100L water applying a minimum of 250mL 4FARMERS TAKE UP per hectare .

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION .