

Bromoxynil 200

Similar to: *Bromicide*® 200

Active Constituent	Bromoxynil as n-octanoyl ester 200g/L		Formulation	EC
Typical Situations	Wheat, oats, barley, triticale, pasture and other (see registration section).			
Chemical Group	C	Mode of Action	Foliar uptake, poor translocation – disrupts photosynthesis and destroys cells.	
Indicative Price ex GST	\$10.50/L	Typical Pack Size	20L, 110L, 1000L	
Poison Schedule	6	Dangerous Goods Class	-	UN -
Packing Code-				
Withholding Period	Harvest	N/A		
	Grazing	Do not graze or cut for hay for 8 weeks post application.		
Plant Back				
Application Method	Boom spray 50-200L/ha, aerially 30L/ha			Rain fastness
	3 hours			
Efficacy	Apply to actively growing weeds. Good coverage is very important to maximise efficacy. Afternoon or evening application will assist translocation/efficacy.			
Adjuvants	None Required			
Compatibility	Phenoxys, grass selective herbicides, most fungicides and insecticides. (Some antagonisms with MCPA and grass selectives exist – see label)			
Incompatibility	Fenoxaprop-P-ethyl products are antagonistic. Incompatible with Glyphosate and Bipyridyls (Paraquat and Diquat).			
Water Quality				
Time to Effects and Symptoms	4-7 days - leaves turn chlorotic and desiccation occurs.			
4F Broadacre Registrations	Wheat, oats, barley, triticale, lucerne and pasture (extract listed).			
4F Other Registrations	Roadsides, Rights of Way and Turf (see label).			
Similar Product Registrations	<i>Bromicide</i> ® 200 is registered for Clover seed crops and fallow situations.			
Situation	Target Weed/s	Rate/ha	Comments	
Cereals (3 leaf to beginning of stem elongation Z13-30)	Amsinckia, Capeweed, Saffron thistle, Variegated thistle	1.4L	Effective on weeds up to 4 leaf or 35mm diameter.	
	Mexican Poppy, Turnip weed, Wild Radish, Wild Mustard, Wild Turnip, Paterson's Curse, Fumitories	2L	Effective on weeds up to the 4 leaf stage or 50mm diameter.	
Pastures	Doublegee, Wild Radish (up to 4 leaf)	1.5L	Apply May-June.	
	Matricaria (up to 3 leaf)	750ml		
Comments				
<p>Do not apply on days where temperatures above 20°C occur or when these temperatures are expected within several days of treatment; may result in seedling mortality and established plants may be damaged. Wheat varieties Shortim and Olympic show susceptibility to Bromoxynil 200. Clovers are moderately tolerant however medics and lucerne will suffer substantial damage at low rates and be killed at high rates.</p>				