

Omethoate 290

Similar to: *Le-mat*[®]

Active Constituent	Omethoate 290g/L		Formulation SL	
Typical Situations	Barrier spray (firebreaks and fence lines).			
Insects Controlled	Red Legged Earth Mite.			
Chemical Group	1B	Mode of Action	Short-term systemic in plants, contact and ingestion with some surface residual activity.	
Indicative Price ex GST	\$23.00/L	Typical Pack Size	10L	
Poison Schedule	6	Dangerous Goods Class	6.1/3	UN 3017 Packing Code III
Withholding Period	Harvest	Not required when used as directed.		
	Grazing	Not required when used as directed.		
Application Method	Boom spray or offset jet 50-100L/ha			Rain fastness 1 hour
Efficacy	Ensure thorough coverage of both soil and weeds in the barrier area (via appropriate jet size and water rates).			
Compatibility	Compatible with most commonly used herbicides (conduct a jar test if unsure).			
Incompatibility	Not compatible with alkaline mixtures (e.g. Bordeaux mixture and lime).			
Water Quality				
4F Broadacre Registrations	N/A			
4F Other Registrations	Barrier spray (firebreaks and fence lines).			
Similar Product Registrations	As above.			
Situation	Target Insect/s	Rate/ha	Comments	
Barriers	Red Legged Earth Mite	300mL	To prevent an invasion, spray bare soil around the crop/pasture along the fence line.	
Comments				
CARE: Damaging to bees - to minimise impact on foraging bees, apply early morning or late evening. Omethoate can provide a residual of 1-2 weeks for the control of Red Legged Earth Mite.				