

Triadimefon 500 Dry

Similar to: *Unique to 4Farmers*

Active Constituent	Triadimefon 500g/kg		Formulation WP	
Typical Situations	Wheat and barley.			
Diseases Controlled	Barley Powdery Mildew, Wheat Leaf and Stripe Rust and Take-all (suppression).			
Chemical Group	3	Mode of Action	Upwardly systemic in plant – provides protectant, curative and eradivative activity.	
Price ex GST	\$19.80/kg	Typical Pack Size	20kg	
Poison Schedule	6	Dangerous Goods Class	-	UN - Packing Code -
Withholding Period	Harvest	Do not harvest for 3 months post application.		
	Grazing	Do not graze or cut treated foliage for stock food.		
Application Method	Mixed dry with fertiliser and applied in-furrow		Rain fastness	N/A
Efficacy	Apply product evenly onto the fertiliser. Ideally the powder should be metered via an applicator into an auger or elevator.			
Adjuvants	None Required			
Compatibility				
Incompatibility				
Water Quality				
4F Broadacre Registrations	Wheat and barley.			
4F Other Registrations	N/A			
Similar Product Registrations	N/A			
Situation	Target Disease/s	Rate/ha	Comments	
Barley	Powdery Mildew	200g	Use when disease risk is high, and monitor crops from 10-12 weeks for follow up control requirements. Up to 16 weeks control not uncommon.	
Wheat	Leaf Rust, Stripe Rust Suppression: Take All			
Comments				
Used as a fertilizer treatment, Triadimefon provides cost effective protectant activity for the above diseases; when residual activity declines and/or where disease persists, another product should be used for curative activity.				
CARE: May impact coleoptile length in some circumstances; ensure adequate seed/fertiliser separation, particularly when sown into pre-emergent herbicide treated soil.				
Dangerous to aquatic organisms and soil fungi (temporary).				