

# Triclopyr 755

Similar to: *Garlon*®

<b>Active Constituent</b>	Triclopyr present as Butoxyethyl ester 755g/L		<b>Formulation</b>	EC
<b>Typical Situations</b>	Fallow/summer weed control, non-crop and Sorghum.			
<b>Chemical Group</b>	I	<b>Mode of Action</b>	Foliar and some root uptake, readily translocated – causes unregulated growth.	
<b>Indicative Price ex GST</b>	\$16.00/L	<b>Typical Pack Size</b>	20L, 110, 1000L	
<b>Poison Schedule</b>	6	<b>Dangerous Goods Class</b>	-	<b>UN</b> - <b>Packing Code-</b>
<b>Withholding Period</b>	<b>Harvest</b> Not required when used as directed. <b>Grazing</b> Not required when used as directed.			
<b>Plant Back</b>	Wheat, barley, sorghum, maize, chickpeas, soybeans & sunflowers - 7 days. Cotton - 14 days.			
<b>Application Method</b>	Boom spray 50L/ha, aerially 35L/ha			<b>Rain fastness</b> 1 hour
<b>Efficacy</b>	Do not spray on stressed weeds.			
<b>Adjuvants</b>	The addition of crop oil (e.g. Turbo Charge) can assist efficacy on difficult to control weeds (do not use crop oil when mixing with Glyphosate).			
<b>Compatibility</b>	Glyphosate, Picloram, Clopyralid, Fluroxypyr and 2, 4-D, and Chlorpyrifos.			
<b>Incompatibility</b>				
<b>Water Quality</b>	Hard water (>200ppm calcium carbonate) may reduce efficacy.			
<b>Time to Effects and Symptoms</b>	3-10 days – twisting stems, cupping leaves and chlorosis at growing points. 10-56 days – wilting and death.			
<b>4F Broadacre Registrations</b>	Fallow/summer weed control (extract listed).			
<b>4F Other Registrations</b>				
<b>Similar Product Registrations</b>	4Farmers Triclopyr 600, other fallow weeds, non-crop, commercial, industrial, rights of way and pasture (extract listed).			
<b>Situation</b>	<b>Target Weed/s</b>	<b>Rate/ha</b>	<b>Comments</b>	
Fallow, Stubble, Firebreak	Camel (afghan) Melon	Up to 20cm in Diameter	95ml	Use with Turbo Charge at 500ml/100L (though not if mixed with Glyphosate).
		Runners from 20 to 40cm in Diameter	130ml	
	Prickly Paddy Melon	Up to 20cm in Diameter	65ml	Use with Turbo Charge at 500ml/100L (though not if mixed with Glyphosate).
		Runners from 20 to 40cm in Diameter	130ml	
		65-130ml + 750ml - 1.5L Glyphosate	Do not add crop oil. Add ammonium sulphate and non-ionic surfactant.	
		65-130ml + 750ml - 1.5L Glyphosate	Do not add crop oil. Add ammonium sulphate and non-ionic surfactant.	
<b>Comments</b>				
Triclopyr provides effective control of summer melons, and is often used as a Glyphosate spike.				
<b>Do not</b> apply to weeds that are under stress, or not actively growing.				
<b>Do not</b> apply to wet foliage.				
<b>Do not</b> clear/burn woody weeds for at least 6 months post treatment.				
<b>CAUTION:</b> Broadleaf plants are highly susceptible – follow label recommendations to avoid off-target impacts.				
Remember to follow label restrictions of tank mix partners.				