

# Fenamiphos 400

Similar to: *Nemacur*<sup>®</sup>

<b>Active Constituent</b>	Fenamiphos 400g/L		<b>Formulation</b> EC	
<b>Typical Situations</b>	Banana and Aloe Vera planting material.			
<b>Insects Controlled</b>	Soil borne plant parasitic nematodes.			
<b>Chemical Group</b>	1B	<b>Mode of Action</b>	Systemic in plants, contact and ingestion activity.	
<b>Price ex GST</b>	\$30.15/L	<b>Typical Pack Size</b>	20L	
<b>Poison Schedule</b>	7	<b>Dangerous Goods Class</b>	6.1	<b>UN</b> 3018 <b>Packing Code</b> II
<b>Withholding Period</b>	<b>Harvest</b>	Not required when used as directed.		
	<b>Grazing</b>	Not required when used as directed.		
<b>Application Method</b>	Dipping		<b>Rain fastness</b>	N/A
<b>Efficacy</b>	Follow label recommendations.			
<b>Compatibility</b>	-			
<b>Incompatibility</b>	-			
<b>Water Quality</b>	-			
<b>4F Broadacre Registrations</b>	-			
<b>4F Other Registrations</b>	Banana and Aloe Vera planting material.			
<b>Similar Product Registrations</b>	As above.			
<b>Situation</b>	<b>Target Insect/s</b>	<b>Rate/100L water</b>	<b>Comments</b>	
Aloe Vera (planting material)	Soil borne plant parasitic nematodes	1L	Allow pups to air dry or cure for 48 hours prior to treatment. Immerse pups in solution for 30 minutes following all safety precautions.	
Banana (planting material)	Soil borne plant parasitic nematodes	100mL	Remove all dead tissue from planting material and dip for 10 minutes. Allow time to drain and dry.	
<b>Comments</b>				
Dip cannot be disposed of on site; by law, mix must be taken to a lawfully recognised disposal facility.				
<b>Do not</b> use this product for at home garden use.				
<b>Do not</b> apply to dry soil or dry grass.				
Wood ducks are particularly susceptible to this chemical use, ensure correct methods of use.				