

# Fipronil 500

Similar to: Cosmos®

<b>Active Constituent</b>	Fipronil 500 g/L		<b>Formulation FS</b>	
<b>Typical Seed</b>	Canola			
<b>Pests Controlled</b>	Red Legged Earth Mite.			
<b>Chemical Group</b>	2B Fungicide	<b>Mode of Action</b>		
<b>Price ex GST</b>	\$400.00	<b>Typical Pack Size</b>		1L
<b>Poison Schedule</b>	6	<b>Dangerous Goods Class</b>	-	<b>UN 2092</b> <b>Packing Code III</b>
<b>Withholding Period</b>	<b>Harvest</b>	Not required when used as directed.		
	<b>Grazing</b>	Canola – do not graze plants grown from treated seed, or cut for stock food, within 9 weeks of sowing.		
<b>Application/Efficacy</b>	Ensure even coverage of seed. Apply with an applicator designed for liquid seed treatments. Calibrate application equipment for the flow rate of grain. Fipronil 500 Insecticidal Seed Treatment should be applied in a total volume of 1.0 L water/100 kg seed.			
<b>Treated Grain</b>	-			
<b>Storage Life</b>	-			
<b>Compatibility</b>	4Farmers Metalaxyl-M 350			
<b>Incompatibility</b>	See above.			
<b>4F Broadacre Registrations</b>	Canola.			
<b>4F Other Registrations</b>	Rice, Sorghum.			
<b>Similar Product Registrations</b>	As above.			
<b>Situation</b>	<b>Target Pest/Disease</b>	<b>Rate/1t seed</b>	<b>Comments</b>	
Canola	Redlegged earth mite (Halotydeus destructor)	400ml/100kg of seed	Ensure thorough coverage of seed. Add 400mL of product to 600mL of water /100kg seed (total volume 1.0L/100kg seed).	
<b>Comments</b>				
Seedling protection will be satisfactory under conditions of low pest pressure (i.e. Less than 1000 per m2). Where greater pest numbers are anticipated or where there are more than 8 mites per plant, alternative or additional treatments will be required.				