

Fipronil 800 WG

Similar to: *Regal*[®] and *Cosmos*[®]

Active Constituent	Fipronil 800g/kg		Formulation WG	
Typical Situations	Pasture, sorghum and various irrigated, viticultural and horticultural crops (see registration section).			
Insects Controlled	Australian Plague Locust, Banana Rust Thrips, Banana Weevil Borer, Cabbage Cluster Caterpillar, Cabbage White Butterfly, Cotton Thrips, Diamondback Moth, Fig Longicorn, Green Mired, Mole Cricket, Mushroom Flies, Spur throated Locust, Sugarcane Weevil, White fringed Weevil, Wingless Grass Hopper and Wireworm.			
Chemical Group	2B	Mode of Action	Systemic in plants, contact and ingestion activity.	
Price ex GST	\$396.00/kg	Typical Pack Size	0.5kg	
Poison Schedule	6	Dangerous Goods Class	6.1	UN 2588 Packing Code III
Withholding Period	Harvest	Do not harvest sorghum for 14 days post application (see label for other crops).		
	Grazing	Do not graze or cut pasture or sorghum for 14 days post application (see label for other crops).		
Application Method	Boom spray at least 50L/ha, aerially at least 20L/ha		Rain fastness	Until dry
Efficacy	Apply to dry foliage and ensure thorough coverage for optimum efficacy.			
Compatibility	-			
Incompatibility	-			
Water Quality	-			
4F Broadacre Registrations	Pasture and sorghum (extract listed).			
4F Other Registrations	Various irrigated, viticultural and horticultural crops (see label).			
Similar Product Registrations	<i>Cosmos</i> [®] (500g/L Fipronil in an SC formulation) is registered for the treatment of canola seed to control Red Legged Earth Mite.			
Situation	Target Insect/s	Rate/ha	Comments	
Pasture and Sorghum	Australian Plague Locust, Spur throated Locust, Wingless Grass Hopper	1.5g	Apply with a minimum of 50L/ha water.	
Comments				
Withhold stock from slaughter for 21 days if they were present at application. Maximum Residue Limits (MRL), Export Slaughter Intervals (ESI) and Export Grazing Intervals (EGI) exist for this product - always check up to date information for Fipronil 800.				