

Proteb

Similar to: *Prosaro*®

Active Constituent	Prothioconazole 210g/L + Tebuconazole 210 g/L		Formulation SC
Typical Situations	For the control of various diseases in wheat, barley, oats, triticale, canola and pyrethrum		
Diseases Controlled	Leaf Rust, Leaf Scald, Net Blotch, Powdery Mildew, Stem Rust, Stripe Rust, Septoria Nodorum		
Chemical Group	3	Mode of Action	Group 3 Fungicide
Price ex GST			Typical Pack Size 5L, 10L, 20L
Poison Schedule	5	Dangerous Goods Class	-
		UN	-
		Packing Code	-
Withholding Period	<p>Canola harvest - not required when used as directed</p> <p>Grazing - do not graze or cut for stock food for 14 days after application</p> <p>Cereals: harvest- do not harvest for 5 weeks after application</p> <p>Grazing - do not graze or cut for stock food for 14 days after application</p> <p>Pyrethrum: harvest and grazing - not required when used as directed</p>		
Application Method	<p>Ground: Apply product using a medium spray quality</p> <p>Aerial: (not pyrethrum) Apply product using a minimum spray volume of 20 L/ha and a medium spray quality</p>		Rain fastness 2-4 hours
Efficacy	-		
Adjuvants	Depending on the disease that is to be treated in the crop, some benefit in efficacy may be gained from addition of an appropriate adjuvant to the spray mixture.		
Compatibility	-		
Incompatibility	-		
Water Quality	-		
4F Broadacre Registrations	Wheat, barley, oats, canola, triticale and pyrethrum.		
4F Other Registrations			
Similar Product Registrations	As Above		
Situation	Target Disease/s	Rate/ha	Comments
Barley (Monitor crops from mid to late tillering).	Powdery mildew	150 - 300mL/ ha	Use the higher rate in higher yielding crops where conditions favour disease development or susceptible varieties are grown.
	Leaf rust		Apply at the first sign of disease development. Monitor and reapply within 14 to 21 days if conditions favour disease development. Use the higher rates (up to 300 ml/ha) where conditions favour severe disease, or disease is established in the lower canopy
Wheat (Monitor crops from early stem elongation)	Stripe Rust	150 - 300ml/ha + adjuvant (refer to Use of Adjuvant)	Use the higher rate (up to 300 ml/ha) in higher yielding crops where conditions favour disease development or susceptible varieties are grown.
	Stem Rust		Continue to monitor crops after application, re-application may be required if conditions favour disease development and initial application is made before the flag leaf has emerged
	Leaf Rust		
Canola (Apply at the 4 to 6 leaf crop stage)	Blackleg	375 - 450ml/ha	Will reduce lodging and stem canker from blackleg. A follow up application may be required at green bud stage in high disease risk situations or where an effective blackleg seed treatment has not been used
Comments:			
Monitor the crop regularly for symptoms of disease. Generally spray at the first sign of disease, although this will depend on factors such as expected weather conditions and the particular crop variety resistance. Refer to Directions for Use for particular disease recommendations. Up to two sprays of 4FARMERS ProTeb SC FUNGICIDE may be applied per season to the crop. Ensure good coverage of all susceptible plant parts.			