

Triadimefon 500 Dry

Similar to: *Unique to 4Farmers*

Active Constituent	Triadimefon 500g/kg		Formulation WP	
Typical Situations	Wheat and barley.			
Diseases Controlled	Barley Powdery Mildew, Wheat Leaf and Stripe Rust and Take-all (suppression).			
Chemical Group	3	Mode of Action	Upwardly systemic in plant – provides protectant, curative and eradivative activity.	
Price ex GST	\$19.80/kg	Typical Pack Size	20kg	
Poison Schedule	6	Dangerous Goods Class	-	UN - Packing Code -
Withholding Period	Harvest	Do not harvest for 3 months post application.		
	Grazing	Do not graze or cut treated foliage for stock food.		
Application Method	Mixed dry with fertiliser and applied in-furrow		Rain fastness	N/A
Efficacy	Apply product evenly onto the fertiliser. Ideally the powder should be metered via an applicator into an auger or elevator.			
Adjuvants	None Required			
Compatibility	-			
Incompatibility	-			
Water Quality	-			
4F Broadacre Registrations	Wheat and barley.			
4F Other Registrations	N/A			
Similar Product Registrations	N/A			
Situation	Target Disease/s	Rate/ha	Comments	
Barley	Powdery Mildew	200g	Use when disease risk is high, and monitor crops from 10-12 weeks for follow up control requirements. Up to 16 weeks control not uncommon.	
Wheat	Leaf Rust, Stripe Rust Suppression: Take All			
Comments				
Used as a fertilizer treatment, Triadimefon provides cost effective protectant activity for the above diseases; when residual activity declines and/or where disease persists, another product should be used for curative activity.				
CARE: May impact coleoptile length in some circumstances; ensure adequate seed/fertiliser separation, particularly when sown into pre-emergent herbicide treated soil.				
Dangerous to aquatic organisms and soil fungi (temporary).				