

Product Name: 4FARMERS IRON CHELATE SNAIL AND SLUG BAIT  
APVMA Approval No: 90221/128109



Label Name:	4FARMERS IRON CHELATE SNAIL AND SLUG BAIT
Signal Headings:	READ SAFETY DIRECTIONS BEFORE OPENING OR USING
Constituent Statements:	60g/kg IRON EDTA COMPLEX
Mode of Action:	
Statement of Claims:	For the Control of Certain Snails and Slugs in a Range of Agriculture and Horticulture Situations as specified in the Directions For Use.
Net Contents:	5 kg - 1000 kg
Restrains:	
Directions for Use:	This section contains file attachment.
Other Limitations:	
Withholding Periods:	DO NOT ALLOW POULTRY OR LIVESTOCK TO FEED ON TREATED AREAS FOR AT LEAST 7 DAYS AFTER APPLICATION.
Trade Advice:	
General Instructions:	GENERAL INSTRUCTIONS: Dead snails will rarely be seen in the open under normal conditions of use as they usually crawl back to their shelters where they die. 4FARMERS

IRON CHELATE SNAIL AND SLUG BAIT is effective at low temperatures, as it does not depend on dehydration for its rapid action.

Resistance Warning:

Precautions:

DO NOT place in heaps. Clean up spilled pellets after filling applicator.

Protections:

PROTECTION OF LIVESTOCK, DOMESTIC ANIMALS, WILDLIFE, FISH, CRUSTACEANS AND THE ENVIRONMENT.  
the product is of low toxicity to mammals, but animals and birds may find this bait attractive. This product is mildly alkaline and may cause some animals, particularly dogs, to vomit if an appreciable quantity (over 1% of body weight) is consumed. If the dog consumes an appreciable amount consult a veterinarian. This product does not contain metaldehyde or methiocarb.  
  
DO NOT contaminate streams, rivers, or waterways with the product or used containers. This product is poisonous to crustacean.

Storage and Disposal:

Store in the closed original container in a dry, cool, well ventilated area out of direct sunlight. Store in a locked place out of the reach of children, domestic or farm animals, poultry and ducks.  
Shake remainder into applicator. Do not dispose of undiluted chemicals on site. Puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

Safety Directions:

MAY IRRITATE THE EYES. AVOID CONTACT WITH EYES. WHEN LOADING OR USING THE PRODUCT WEAR DISPOSABLE DUST FACE MASK COVERING MOUTH AND NOSE. IF PRODUCT GETS IN EYES, WASH IT OUT IMMEDIATELY WITH WATER. WASH HANDS AFTER USE.

First Aid Instructions:

If poisoning occurs contact a doctor or the Poisons Information Centre on Tel No:13 11 26, New Zealand 0800 764 766

First Aid Warnings:

## DIRECTIONS FOR USE

Crop/Situation	Pest	States	Rates	Critical Comments
Pasture (established and seedling) Crop boundaries, Bare ground (including broadacre pre-emergent, pre-establish) Ornamentals Canola Poppies	Grey field slugs	All states	5 to 16kg/ha	Apply at 5-16kg/ha. For most infestation use the lower rate. For heavy infestations use a higher rate. DO NOT heap pellets. When applying to crops, ensure pellets do not become lodged in plant foliage.
Vegetable crops Strawberries Citrus orchards	Common brown(brown) snail White snail Vine snail			
Grapes	Common garden(brown) snail		15kg/ha	
	White snail(vineyard, Italian and conical)	NSW, VIC, TAS, SA, WA		

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORIZED UNDER APPROPRIATE LEGISLATION